




**Algaecide/ Bactericide\*/ Fungicide**  
Labeled for Use on Citrus Crops  
\*Non-public health bacteria



**CITRUS - Grapefruit, Kumquat, Lemon, Lime, Orange, Tangelo, Tangerine**

Disease	Rate per Acre fl. oz.	Minimum retreatment interval (days)	Instructions
Brown Rot	30-70	7	Apply at first indication of rain or first appearance of Brown Rot.
Greasy Spot, Pink Pitting	25.6-64	7	Apply during mid-summer.
Scab	25.6-64	7	Apply shortly before trees begin to flush. Re-apply at 2/3 petal fall. Re-apply 4 weeks later, if necessary.
Melanose	12.8-64	7	Apply 2 times per year (applications must be 7 days apart) before onset of spring and autumn rains.
Canker (Suppression)	12.8-64 **12.6 lbs.	7	Spray flushes 7 to 14 days after shoots begin to grow. Young fruit may require additional applications. Number and timing of applications will be dependent on disease pressure. Under heavy pressure, each new flush of growth should be sprayed. Heavily infected trees should be sprayed with a minimum dosage of 250 ppm with a follow up spray after 7 to 14 days.

\*\*Maximum Annual Load of metallic copper applied per year per acre



**Always read and follow label directions.**

Manufactured by:  
Earth Science Laboratories Inc.  
113 SE 22nd Street, Suite 1  
Bentonville, AR 72712  
800.257.9283  
EarthScienceLabs.com



**Algaecide/ Bactericide\*/ Fungicide**

for Use on Raw, Citrus and Growing Agricultural Commodities

\*Non-public health bacteria

### GENERAL INSTRUCTIONS

ET-F may be applied with any type of application equipment that gives uniform coverage of all foliage, including ground, aerial and low volume sprayers as specified on this label. Equipment used for application should be PVC or 316L stainless steel. ET-F is compatible with most fungal and insecticidal biopesticides when applied at least two (2) days before or after application of the biopesticide.

Phytotoxicity - Although ET-F has been tested on a wide variety of fruits, vegetables and nuts without phytotoxicity, there could be some varieties and cultivars that because of environmental factors and stages of growth could possibly foster systems. The per acre use rate of ET-F is applicable for dilute spraying. Depending on the equipment used and the specific crop, the spray volume applied per acre will differ. Refer to the Minimum Recommended Spray Volume Table. Complete spray coverage is essential to assure optimum performance from ET-F. When treating by aerial application or with low volume application equipment, unless you have had specific previous experience, it is advisable to test for compatibility and tolerance to crop injury prior to full scale commercial utilization.

Consult ET-F label for specific rates and timing of application by crop. Where application rates and intervals are provided in a range (e.g., 2-4 fluid ounces and 7 to 10 days) the higher rates and shorter spray time intervals are recommended when rainfall is heavy and/or disease pressure is high. Use the higher rates for large mature tree crops. The use of a surfactant, such as Cell-U-Wett™ is acceptable for plants having waxy or hairy surfaces. ET-F works via surface contact with the plants and materials being treated. It is important to ensure that all surfaces are thoroughly wetted. ET-F does not produce any visible residue or have a distinct odor. It does have a residual, especially if applied with a surfactant.

### GENERAL CHEMIGATION INSTRUCTIONS

Apply this product only through one or more of the following types of systems: Sprinkler including center pivot, lateral move, end row, side (wheel) roll, traveler, big gun, solid set or hand move: flood (basin); furrow; border or drip (trickle) irrigation and system(s). Do not apply this product through any other type of irrigation systems.

Crop injury, lack of effectiveness, or illegal pesticide residues in the crop can result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service specialists, equipment manufacturers or other experts.

Do not connect an irrigation system (including greenhouse systems) used for pesticide application to a public water system unless the pesticide label prescribed safety device for public water systems are in place.

### FOR SPRAY AND SOIL DRENCH APPLICATIONS

Always spray for total foliage coverage. When re-spraying the rates and severity of the disease vary with unforeseen conditions. However, in the event of severe disease, spraying intervals can be shortened see 'Minimum Retreatment Interval' column for the shortest interval between applications. At times, lower rates can be as effective as higher rates and should be tried first. Usually, preventive programs may be maintained at lower rates. Use of low volume spraying is effective against Botrytis and not effective against established powdery mildew and Xanthomonas infections. Also, applications on actively growing tissue may be more effective than applications on dormant tissue.

MINIMUM RECOMMENDED SPRAY VOLUME (GALLONS) PER ACRE WHEN APPLYING ET-F

GROUND			
CROP	AERIAL	DILUTE	CONCENTRATE*
Citrus	10	125	30
Field Crops	3	20	30
Small Fruits	5	150	30
Tree Crops	10	400	50
Vegetables	3	20	30
Vines	5	150	30

### KEEP OUT OF REACH OF CHILDREN WARNING • AVISO

**IF IN EYES:** Hold eye open and rinse slowly and gently with water for 20 minutes. Remove contact lenses, if present, after first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for advice.

**IF SWALLOWED:** Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by a poison control center or doctor. Do not give anything to an unconscious person.

**IF ON SKIN OR CLOTHING:** Take off contaminated clothing. Rinse skin immediately with plenty of soap and water for 15 to 20 minutes. Call a poison control center or doctor for treatment.

**NOTE TO PHYSICIAN:** Probable mucosal damage may contraindicate the use of gastric lavage.

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact INFOTRAC 1-800-535-5053 for emergency medical treatment.

**SEE ADDITIONAL PRECAUTIONARY STATEMENTS ON LABEL**

# ESLAgri.com